520 Rec'd PCT/PTO 1 2 NOV 1999

ATTORNEY'S DOCKET NUMBER ORM PTO-1390 (Modified) REV 10-95) U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE 99,849-A TRANSMITTAL LETTER TO THE UNITED STATES U.S. APPLICATION NO. (IF KNOWN, SEE 37 CFR DESIGNATED/ELECTED OFFICE (DO/EO/US) 09/423698 CONCERNING A FILING UNDER 35 U.S.C. 371 PRIORITY DATE CLAIMED INTERNATIONAL FILING DATE INTERNATIONAL APPLICATION NO. 14 May 1998 - 1997 14 May 1998 -PCT/FR98/00966 ~ TITLE OF INVENTION MULTIVALENT VACCINE COMPOSITION WIHT MIXED CARRIER  $\diagup$ APPLICANT(S) FOR DO/EO/US LEROY, Odile 🐱 Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information: This is a FIRST submission of items concerning a filing under 35 U.S.C. 371. This is a SECOND or SUBSEQUENT submission of items concerning a filing under 35 U.S.C. 371. 2. This is an express request to begin national examination procedures (35 U.S.C. 371(f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. 371(b) and PCT Articles 22 and 39(1). 3. A proper Demand for International Preliminary Examination was made by the 19th month from the earliest claimed priority date.  $\times$ 4. A copy of the International Application as filed (35 U.S.C. 371 (c) (2)) 5. is transmitted herewith (required only if not transmitted by the International Bureau). has been transmitted by the International Bureau. is not required, as the application was filed in the United States Receiving Office (RO/US). A translation of the International Application into English (35 U.S.C. 371(c)(2)). X A copy of the International Search Report (PCT/ISA/210).  $\boxtimes$ Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371 (c)(3)) are transmitted herewith (required only if not transmitted by the International Bureau). have been transmitted by the International Bureau. b. □ have not been made; however, the time limit for making such amendments has NOT expired. C. have not been made and will not be made. A translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)). An oath or declaration of the inventor(s) (35 U.S.C. 371 (c)(4)). A copy of the International Preliminary Examination Report (PCT/IPEA/409).  $\boxtimes$ =11. A translation of the annexes to the International Preliminary Examination Report under PCT Article 36 <u>-</u>12. (35 U.S.C. 371 (c)(5)). Items 13 to 18 below concern document(s) or information included: An Information Disclosure Statement under 37 CFR 1.97 and 1.98. 13. An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included. 14. A FIRST preliminary amendment.  $\times$ A SECOND or SUBSEQUENT preliminary amendment. A substitute specification. 16. A change of power of attorney and/or address letter. 17. Certificate of Mailing by Express Mail 18. X Other items or information: 19. **Express Mail Certificate; Return Postcard** 

## 420 Rec'd PCT/PTO 1 2 NOV 1999

u.s. application	no.dif known, see 37 cfr /423698	INTERNATIONAL APPLICATION NO. PCT/FR98/00966		ATTORNEY'S DOCKET NUMBER  99,849-A	
20. The following	lowing fees are submitted:.			CALCULATION	S PTO USE ONLY
	L FEE ( 37 CFR 1.492 (a) (1) -	(5)):			
🗀 Search Repo	rt has been prepared by the EPO	or JPO	\$840.00		
	preliminary examination fee pai		\$670.00		
☐ No internation	onal preliminary examination fee onal search fee paid to USPTO (3	paid to USPTO (37 CFR 1.482) 7 CFR 1.445(a)(2))	\$760.00		
Neither interinterinternational	national preliminary examination search fee (37 CFR 1.445(a)(2)	n fee (37 CFR 1.482) nor paid to USPTO	\$970.00		
<ul><li>International and all claim</li></ul>	l preliminary examination fee pains satisfied provisions of PCT Am	d to USPTO (37 CFR 1.482) ticle 33(2)-(4)	\$96.00		
l	ENTER APPROPRI	ATE BASIC FEE AM	OUNT =	\$970.00	
Surcharge of \$130.0 months from the ea	00 for furnishing the oath or declaritiest claimed priority date (37 C	aration later than 20 FR 1.492 (e)).		\$130.00	
CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE		
Total claims	23 - 20 =	3	x \$18.00	\$54.00	
Independent claims	2 - 3=	0	x \$78.00	\$0.00	ļ
Multiple Depender	nt Claims (check if applicable).			\$0.00	
		ABOVE CALCULAT		\$1,154.00	
Reduction of 1/2 for must also be filed (	or filing by small entity, if application (Note 37 CFR 1.9, 1.27, 1.28) (ch	able. Verified Small Entity State neck if applicable).	tement	\$0.00	
The state of the s		SUB'	TOTAL =	\$1,154.00	
Processing fee of \$ months from the ea	130.00 for furnishing the English rliest claimed priority date (37 C	translation later than $\Box$ 2 CFR 1.492 (f)).	0 □ 30 +	\$0.00	
		TOTAL NATIONAL	L FEE =	\$1,154.00	
Eee for recording to accompanied by an	he enclosed assignment (37 CFR appropriate cover sheet (37 CFR	1.21(h)). The assignment must 3.28, 3.31) (check if applicab	be le).	\$0.00	
		TOTAL FEES ENCI	LOSED =	\$1,154.00	
				Amount to be: refunded	\$
<u>.</u>				charged	\$
	n the amount of \$1,154.00 arge my Deposit Account No.	to cover the above fees is en		to cover the ab	ove fees.
A duplica	tte copy of this sheet is enclosed.				
	nissioner is hereby authorized to take Account No. 13-2490	charge any fees which may be re A duplicate copy of this sheet		any overpayment	
NOTE: Where at 1.137(a) or (b)) m	n appropriate time limit under ust be filed and granted to rest	37 CFR 1.494 or 1.495 has not ore the application to pending	been met, a peti status.	tion to revive (37 C	FR
SEND ALL CORE	RESPONDENCE TO:	<i>,</i>	W. O. C	1.1-M 0	
Michael S. Green	nfield		SIGNATURE		
Reg. No. 37,142	non Hulbort & Boroboff			VI	
300 South Wack	nen Hulbert & Berghoff er Drive		Michael S. (	Greenfield	
Chicago, Illinois	60606		NAME		
Phone: 312-913-			37,142		
Facsimile: 312-9	13-0002		REGISTRAT	ION NUMBER	
			Navambar 1	2 1000	
M			November 1	14, 1777	
			DATE		

# 420 Rec'd PCT/PTO 1 2 NOV 1999

## IN THE UNITED STATES PATENT AND TRADEMARK OFFICE (Case No. 99,849-A)

In the Appli	cation of:	)
	Odile Leroy	) Examiner: TBA
Serial No.	U.S. National Phase of PCT/FR98/00966	) Group Art Unit: TBA
Filed:	November 12, 1999	)
Title:	Multivalent Vaccine Composition With mixed Carrier	) ) )

#### **PRELIMINARY AMENDMENT**

Asst. Commissioner for Patents Washington, D.C. 20231

Dear Sir:

Please consider the following amendments and remarks before calculation of the filing fee.

#### In the Claims:

- 4. (Amended) The [Composition] composition according to [Claim1, 2 or 3,] Claim 1 in which "n" is a number equal to or greater than 6.
- 6. (Amended) The [Composition] composition according to [one of Claims 1 to 5,]

  Claim 1 in which the carrier proteins P1 to Pn are independently selected from a group consisting of two carrier proteins A1 and A2.
- 8. (Amended) The [Composition] composition according to [one of Claims 1 to 7,]

  Claim 1 in which at least one of the carrier proteins P1 to Pn is the diphtheria toxoid (Dt) and at least one of the carrier proteins P1 to Pn is the tetanus toxoid (Tt).

10. (Amended) <u>The [Composition] composition</u> according to [Claim 8 or 9,] <u>Claim 8</u> in which the quantity of Dt is less than or equal to 200 μg/dose.

11. (Amended) The [Composition] composition according to [Claim 8, 9 or 10,] Claim 8 in which the quantity of Tt is less than or equal to 50µg/dose.

19. (Amended) <u>The</u> [Composition] <u>composition</u> according to [Claims 16, 17, or 18,] <u>Claim 16</u> in which "n" is a number equal to or greater than 6.

21. (Amended) The [Composition] composition according to [one of Claims 16 to 20,] Claim 16 in which the polysaccharides S1 to Sn are of bacterial origin.

#### **REMARKS**

No new matter has been added to the application by way of these Amendments.

Respectfully submitted,

Date: November 12, 1999

By: Michael S. Greenfield

Reg. No. 37,142

15

20

25

30

35

WO 98/51339

more pathogenic agent.

## Multivalent vaccine composition wi

The subject of the present invention is pharmaceutical composition intended for the treatment or prevention of a number of infections caused by pathogenic agents such as bacteria, which comprises, as immunogenic agent, polysaccharides derived from one or

Bacteria as well as fungi such as incorporate polysaccharides into their surface structure. Thus, the great majority of bacteria are coated with an exudate of a polysaccharide nature which is attached to the bacterium more or less firmly but which is strictly speaking not an envelope. This exudate is called glycocalyx or capsule. Moreover, the outer membrane of Gram-negative bacteria, consists, inter alia, of lipopolysaccharide (LPS). Finally, polysaccharides are also found in the wall of fungi. These polysaccharides are in fact surface antigens which induce an immune response in an infected mammal.

Such polysaccharides are produced on the basis of units in which the constituents and the bonds are defined and which are characteristic of the bacterial or fungal species considered. These repeating units contain the epitopes, that is to say the structures which determine antigenicity.

The polysaccharides οf pathogenic organisms are reputed to be good vaccine agents. As they are, they are effective in adults and children over two years. On the other hand, in breast-feeding infants, some are only slightly or not immunogenic and do not induce any immune response. It is possible to overcome this problem by coupling, via bonding, the polysaccharides to a so-called carrier protein such as diphtheria or tetanus toxoid so as to obtain a polysaccharide-carrier protein conjugate.

same vaccine composition mav several conjugates. Indeed, the trend is to combine several vaccinal agents intended to prevent or to treat

15

20

25

30

infections induced by pathogenic agents from various species, this being, inter alia, in order to limit the number of administrations during the life of individual. Furthermore, within the same species, there may be several serogroups/serotypes which are widely represented regionally or world-wide, It is this recalled that a serogroup/serotype is characterized, inter alia, by the nature of the capsule polysaccharide that polysaccharides of various serogroups generally đo not exhibit immunological reactivity. In this case, it may therefore be necessary to combine the polysaccharides obtained from various serogroups in order to effectively combat an infection caused by one and the same species.

Thus, this is for example the case when it is sought to vaccinate against Streptococcus pneumoniae infections. Pneumococcal infections are a real public health problem especially since they are found in the severe forms of pneumonia, septicaemia and meningitis. In industrialized countries, they affect each year 30 to 100 per 100,000 children under three years. The mortality rate in cases of bacteraemia and meningitis is 15 to 30% whereas 5% of children die of pneumonia.

A study carried out in Finland from 1985 to 1989 shows that 90% of invasive infections are caused groups of serogroups/serotypes. Serogroups/ serotypes 14, 6 and 19 are responsible for 54% of cases, serotype 14 being predominant in children under two years. Other pneumococci frequently isolated belong to serogroups 7, 18 and 23; yet others, more rare, belong to serogroups/serotypes 9 and 4. A similar has distribution been demonstrated in industrialized countries, in particular in the United States.

Moreover, Streptococcus pneumoniae is responsible for a number of otitis infections which are more benign but very common. The number of children that have had an otitis infection before the age of six is evaluated at about 75% and the number of otitis

15

infections caused by pneumococcus at 30 to 50%. In developed countries, otitis infections caused by pneumococcus are due to serogroup 19 in 25% of cases, followed by serogroups/serotypes 23 (13%), 6 and 14 (12%), 9 and 18 (4%) and 4 and 1 (2%).

A pneumococcal vaccine containing the polysaccharides of 23 serotypes is already commercially available. This vaccine makes it possible to effectively combat invasive infections in adults and has a transient action in children over seven months.

The capsular polysaccharides of pneumococci are T-independent antigens, i.e. they can induce antibodies, preferably of the IgM type, without the help of T cells and are not capable of promoting a booster response of the IgG type. When they are coupled to a carrier protein, these polysaccharides then prove capable of inducing a T-dependent response, most particularly in neonates and should provide long-term protection.

studies have been carried out 20 Clinical Finland and Israel with pneumococcal vaccines having four valencies containing conjugates 6B, 9V, 18C and 23F in which the polysaccharide was coupled either to Dt or to Tt. The doses were 1, 3 or polysaccharide per valency. Each of these formulations 25 administered simultaneously with an Haemophilus vaccine (polyribitolphosphate coupled to Tt; Act-HIB marketed by Pasteur Mérieux Connaught) and an anti-diphtheria, tetanus, whooping cough vaccine (for Finland, D.T.Coq marketed by KTL). Furthermore, 30 three administrations were carried accompanied or not by simultaneous administration of an oral or injectable polio vaccine. They were repeated twice at a few weeks interval, and then once, one year after the first immunization. 35

The results of these studies as reported in the table below have made it possible to demonstrate an effect of negative interference of the diphtheria and

tetanus toxoid load on the induction of anti-HiB antibodies, after the last immunization.

Finnish study		study	Anti-HiB antibody in µg/ml	
Placebo			11.00	
Tetra	valen	t pneumo		
Tt:	1	ħã	10.1	
	3	μg	7.18	
	10	μg	4.11	
Dt:	1	μg	11.5	
	3	μg	12.5	
	.10	μg	7.18	

Israeli study	Anti-HiB antibody in
	μg/ml
Placebo	6.62
Tetravalent pnew	по
Tt: 3 µg	2.81
Dt: 3 µg	4.62

5

A similar interference effect was observed during a clinical study in Iceland in which breast-feeding infants received Pro HIBit (PRP coupled to Dt; Connaught) in place of Act-HIB.

10 More generally, it is predicted regardless of the vaccine based on conjugated polysaccharides, a maximum load of Dt and of Tt or of any other protein exists in the conjugated vaccine or in the association or combination of vaccine administered above which the immune response against polysaccharides conjugated with this protein may be reduced. In order to overcome the problem which the phenomenon of negative interference constitutes in multivalent vaccines composed of polysaccharide conjugates, the 20 present application proposes to use not one but at least two carrier proteins so that the maximum load of each of the carrier proteins is not reached.

Accordingly, the subject of the invention is a composition comprising "n" conjugates C1 to Cn, each conjugate being composed (i) of a polysaccharide, in particular a polysaccharide derived from a Streptococcus pneumoniae serotype/serogroup S1 respectively, and (ii) of a carrier protein P1 to Pn respectively, "n" being a number equal to or greater than 2; in which composition the polysaccharides S1 to Sn are identical or different and in which the carrier proteins P1 to Pn are selected independently from a group consisting of "m" carrier proteins Al to Am, "m" being a number equal to or greater than 2, provided that at least one of the carrier proteins P1 to Pn is different from the others.

15 According to another aspect, the subject of the invention is also a composition which comprises "n" conjugates C1 to Cn, each conjugate being composed (i) of a polysaccharide S1 to Sn respectively and (ii) a carrier protein P1 to Pn respectively, "n" being a number equal to or greater than 2; in which composition 20 the polysaccharides S1 to Sn are identical or different and in which the carrier proteins P1 to Pn are selected independently from a group consisting of diphtheria (Dt) and tetanus (Tt) toxoids, provided that at least one of the carrier proteins P1 to Pn is different from 25 the others; and which is characterized in that the quantity of Dt and Tt is respectively less than or to 200 50 μg/dose. and In other words, composition according to the invention comprises one or 30 more polysaccharide conjugates in which polysaccharide is coupled to the diphtheria toxoid (Dt) and one or more polysaccharide conjugates in which the polysaccharide is coupled to the tetanus toxoid (Tt) and is characterized in that the quantity of Dt and Tt is respectively 35 less than or equal 200 to 50 µg/dose.

By way of illustration, the following compositions are envisaged:

20

25

30

35

- (i) A composition containing at least three conjugates C1, C2, C3, ... Cn, of formulas S1-P1, S2-P2, S3-P3, .... Sn-Pn, with: S1 to Sn identical to each other and P1 to Pn all different from each other;
- 5 (ii) A composition containing at least three conjugates C1, C2, C3, ... Cn, of formulas S1-P1, S2-P2, S3-P3, .... Sn-Pn, with: S1 to Sn all different from each other, P1 and P2 identical to each other, P3 to Pn identical to each other and P1 and P2 different from P3 to Pn, and
  - (iii) A composition containing at least three conjugates C1, C2, C3, ... Cn, of formulas S1-P1, S2-P2, S3-P3, .... Sn-Pn, with: S1 and S2 identical to each other, S3 to Sn identical to each other, S1 and S2 different from S3 to Sn, P1 and P3 identical to each other, P2 to Pn, excluding P3, identical to each other and P1 and P3 different from P2 to Pn (-P3).

Thus. for the purposes of the present conjugates invention, the C1 to Cn, which necessarily all different from each other, may be so in pairs either through their polysaccharide, or through their carrier protein or through their polysaccharide and their carrier protein. According to a specific embodiment, the polysaccharides used are all different from each other.

The number "n" of conjugates present in composition according to the invention is equal to or greater than 2, and may in particular be equal to or greater than 3, 4, 6, 8, 10, 11, 12, 15 or 20. general, this number "n" may be determined by persons skilled in the art as a function of a number of criteria in particular linked to the very nature of the composition, to the objectives which this composition should make it possible to achieve and population for whom this composition is intended. For example, in the case of a composition intended for treating or preventing pneumococcal infections breast-feeding infants, it is considered that such a composition, in order to offer a good level of

10

35

protection and world-wide protection, should contain at least 8, preferably at least 10, most preferably at least 11 valencies which may be represented by at least 11 conjugates or more.

"Polysaccharide" is understood to mean a polymer consisting of a plurality of saccharide repeating units, especially of more than four repeating units, regardless of the length of the saccharide chain and regardless of the average molecular weight of the polysaccharide. This term covers in particular that of oligosaccharide.

"Conjugate" is understood to mean a compound in which a polysaccharide is covalently linked to a carrier protein.

15 Thus, as previously stated, a composition according to the invention should use at least two carrier proteins. These carrier proteins may be chosen from all those commonly used in the field of vaccines. They may be in particular the diphtheria toxoid (Dt), the tetanus toxoid (Tt), the non-toxic mutant form 20 CRM197 of the diphtheria toxin and the outer membrane protein type 1 (OMP1) of Neisseria meningitidis or any variant, analogue or fragment of the latter which has preserved the carrier capacity. The methods which make it possible to obtain these proteins in purified form 25 are well known to persons skilled in the art. The terms "protein" as used in the present application applies to any amino acid chain, regardless of the length of the chain. In particular, this term covers the notion of 30 peptide.

In general, the group of proteins Al to Am from which the carrier proteins P1 to Pn are independently selected therefore represents all the proteins having a carrier effect. For their personal needs, skilled in the art may agree that their choice would be limited to a defined number of proteins consequently, they can define the group which they will use to make their selection on the basis of a number "m" of components equal to or greater than 2 and at

10

15

20

25

30

35

most equal to "n", "n" being as defined above. In particular, persons skilled in the art can determine the minimum number of different carrier proteins which is necessary in order to avoid the phenomenon of interference. To do this, they will take into account the maximum load that should not be exceeded for each of the carrier proteins. "Maximum load" refers to the quantity of carrier protein above which a reduced immune response is observed against one polysaccharides compared with a corresponding monovalent composition (conjugate taken separately).

In particular, as regards the diphtheria toxoid and the tetanus toxoid, it is estimated that, advantageously, the quantity of these proteins present in a dose of a composition according to the invention should not exceed 200 and 50 µg respectively, such a dose being envisaged for administration in a mammal, preferably a human. Preferably, the Dt load is less than or equal to 150, 120 or 100 µg, most preferably 80 or 60 µg. Preferably, the Tt load is less than or equal to 35 or 25 µg, most preferably 20 or 10 µg.

Thus, it may be accepted that for a composition using only two different carrier proteins, the selection of these proteins will be made from a group consisting of proteins A1 and A2. Preferably, A1 and A2 may be diphtheria toxoid (Dt) and tetanus toxoid (Tt) respectively or vice versa.

According to specific a embodiment, composition using only two different carrier proteins is characterized by a balanced distribution of the number of polysaccharides conjugated with the first carrier protein and of the number of polysaccharides conjugated with the second carrier protein. example, when "n" is an even number, "n"/2 carrier proteins P1 to Pn are A1 and "n"/2 carrier proteins P1 to Pn are A2 or when "n" is an odd number, ("n"+1)/2 carrier proteins P1 to Pn are A1 and ("n"-1)/2 carrier proteins P1 to Pn are A2.

A polysaccharide useful for the purposes of the present invention may be in particular a polysaccharide of bacterial or fungal origin. It may be in particular a polysaccharide from Streptococcus e.g. Streptococcus pneumoniae, Staphylococcus, Klebsiella, Salmonella e.g. Salmonella typhi, Escherichia, Shigella, Neisseria e.g. Neisseria meningitidis, Haemophilus e.g. Haemophilus influenzae, Moraxella, Vibrio cholerae or Mycobacterium tuberculosis.

10 In a composition according to the invention, the polysaccharides may be derived from different species or alternatively may all be derived from the same species, e.g. from the same bacterial species, possibly of different serogroups/serotypes. In order to 15 illustrate this last possibility, there may mentioned a composition according to the invention intended to vaccinate against pneumococcal infections, which contains at least 8 valencies, preferably 10 or 11 valencies chosen from serotypes 1, 3, 4, 5, 6B, 7F, 9V, 14, 18C, 19F and 23F. 20

Thus, a composition constituting a pneumococcal vaccine advantageously comprises 10 or 11 valencies, e.g. represented by 10 or 11 conjugates in which the polysaccharides are all different from each other and are derived (have as origin) serotypes/serogroups chosen from serotypes 1, 3, 4, 5, 6B, 7F, 9V, 14, 18C, 19F and 23F of S. pneumoniae. It may be in particular a composition which comprises 10 or 11 conjugates selected from:

- 30 serotype 1 polysaccharide coupled to Tt or to Dt;
  - serotype 3 polysaccharide coupled to Dt;
  - serotype 4 polysaccharide coupled to Tt;
  - serotype 5 polysaccharide coupled to Tt or to Dt;
  - serotype 6B polysaccharide coupled to Dt;
- 35 serotype 7F polysaccharide coupled to Tt or to Dt;
  - serotype 9V polysaccharide coupled to Tt;
  - serotype 14 polysaccharide coupled to Dt;
  - serotype 18C polysaccharide coupled to Dt;
  - serotype 19F polysaccharide coupled to Tt; and

10

30

35

serotype 23F polysaccharide coupled to Tt.

Under another aspect, a composition constituting a pneumococcal vaccine may comprise 10 or 11 valencies represented by 12 to 22, especially 12 to 15 conjugates, in which the polysaccharides are derived from the serotypes chosen from serotypes 1, 3, 4, 5, 6B, 7F, 9V, 14, 18C, 19F and 23F, in which composition conjugates of the same valency differ from each other in the carrier protein. It may be in particular a composition which comprises:

- serotype 1 polysaccharide coupled to Tt;
- serotype 3 polysaccharide coupled to Dt;
- serotype 4 polysaccharide coupled to Tt;
- serotype 5 polysaccharide coupled to Tt;
- 15 serotype 6B polysaccharide coupled to Dt;
  - serotype 6B polysaccharide coupled to Tt;
    - serotype 7F polysaccharide coupled to Tt;
    - serotype 9V polysaccharide coupled to Tt;
    - serotype 9V polysaccharide coupled to Dt;
- 20 serotype 14 polysaccharide coupled to Dt;
  - serotype 18C polysaccharide coupled to Dt;
  - serotype 18C polysaccharide coupled to Tt;
  - serotype 19F polysaccharide coupled to Tt;
  - serotype 23F polysaccharide coupled to Tt; and
- 25 serotype 23F polysaccharide coupled to Dt.

Such a polysaccharide may be advantageously the microorganism according extracted from methods and purified likewise. conventional This polysaccharide may be used in the crude form after extraction/purification. Alternatively, it fragmented in order to obtain a polysaccharide having an average molecular weight less than that of the polysaccharide originally extracted. Α particularly is advantageous fermentation method described WO 93/7178 which is incorporated by way of reference.

A conjugate in which a polysaccharide is coupled by covalent bonding to a carrier protein may be obtained according to conventional methods well known to persons skilled in the art. It may make use of a

linker or a spacer to carry out the conjugation. Depending on the mode of conjugation used, the conjugate resulting therefrom may be a conjugate in which the polysaccharide is linked to the protein by a single chemical function (sun or neoglycoconjugate type), or by several functions (random coil type). It is within the capability of persons skilled in the art to determine the most appropriate mode of conjugation as a function of the nature of the polysaccharide and more particularly of the chemical groups carried by the polysaccharide which may be used during the conjugation reaction.

A composition according to the invention may be manufactured conventionally. In particular, it may be formulated with a pharmaceutically acceptable diluent 15 or vehicle, e.g. a saline solution. water or addition, the composition may contain customary ingredients such buffer, as a a preservative or stabilizer, an adjuvant such as an aluminum compound, e.g. an aluminium hydroxide, an aluminium phosphate or 20 an aluminium hydroxyphosphate, and, where appropriate, a lyophilization excipient. In general, these products may be selected as a function of the mode and route of administration and based on standard pharmaceutical practices. Appropriate carriers or diluents as well as 25 what is essential for the preparation of a pharmacomposition are described in Remington's ceutical Pharmaceutical Sciences, a standard reference book in this field.

A composition according to the invention may be administered by any conventional route which is used in the field of vaccines, in particular by the systemic, i.e. parenteral, route, e.g. by the subcutaneous, intramuscular, intradermal or intravenous route, or by the mucosal route, e.g. by the oral or nasal route.

The administration may take place in a single dose or in a dose repeated once or several times, e.g. once, twice or three times, after a certain interval of time. The appropriate dosage will vary as a function of

various parameters, for example the number of valencies contained in the composition, the nature of the polysaccharide(s) used or the mode of administration. As a guide, it is indicated that good results may be obtained with per valency, a polysaccharide dose of 0.5 to 100  $\mu$ g, preferably of 1 to 50  $\mu$ g, most preferably of 1 to 10  $\mu$ g. A dose of the composition according to the invention may be advantageously in a volume of 0.1 to 2 ml.

10 There are presented below, by way of example, various pneumococcal vaccines having multiple valencies, the valencies being chosen from serotypes 1, 5, 6B, 7F, 9V, 14, 18C, 19F and 23F. polysaccharides derived from these serotypes 15 fragmented according to the method described in WO 93/7178. The polysaccharides coupled to Tt (except the type 1 polysaccharide) were coupled according to the conjugation method described in WO 93/7178. Briefly, a polysaccharide is subjected to reductive amination in the presence of sodium cyanoborohydride in order to 20 link a molecule of diaminohexane to a terminal reducing The polysaccharide thus derived activated by a succinimide group using disuccinimidyl suberate (DSS). The polysaccharide thus activated is 25 reacted directly with the carrier protein. The serotype 1 polysaccharide was coupled Dt to Tt [sic] according to the conjugation method described in US patent No. 5,204,098 which is incorporated by way of reference. The other polysaccharides coupled to Dt were coupled as follows; hydrazide groups were incorporated onto the 30 polysaccharide by reacting the polysaccharide with an excess of adipic acid dihydrazide (ADH) in the presence of ethyl dimethyl amino propyl carbodiimide (EDAC) and sodium cyanoborohydride (for all the types except 3) or simply in the presence of sodium cyanoborohydride (for 35 type 3). The polysaccharide thus derived is reacted with the carrier protein in the presence of EDAC. The experimental conditions were controlled so as to obtain conjugates in which the quantity of protein is between

one and four times, preferably twice, the value of the quantity of polysaccharide. Thus, for a dose, 3  $\mu g$  of a particular polysaccharide are coupled to about 6  $\mu g$  of Dt and 1  $\mu g$  of a particular polysaccharide is coupled to about 2  $\mu g$  of Tt.

The Dt and Tt used were prepared by detoxification with formaldehyde starting with toxins extracted respectively from *Corynebacterium diphtheriae* and *Clostridium tetani*.

The formulations contain a phosphate buffer (0.475 mg of PO<sub>4</sub><sup>2-</sup> ion per dose) and sodium chloride (4.5 mg per dose) and may be supplemented with aluminium hydroxide (alum) adjuvant (300 µg of Al<sup>3+</sup> ion per dose) and contain a preservative such as phenoxyethanol formalin. A dose is in the volume of 0.5 ml.

EXAMPLE 1: Octavalent formulation

	Polysaccharide		
Carrier protein	Serotype	Quantity per single	
		dose	
Dt	3	3 µg	
Tt	4	1 μg	
Dt	6B	10 μg	
Tt	9V	1 μg	
Dt	14	3 µg	
Dt	18C	3 μg	
Tt	19F	1 μg	
Tt	23F	1 μg	

5

EXAMPLE 2: Formulations F3, F4 and F3bis containing 11 valencies

•	Polysaccharide			
	Serogroup/	Qty per	Qty per	Qty per
Carrier	Serotype	single dose	single dose	single dose
Protein		of formu-	of formu-	of formu-
		lation F3	lation F4	lation F3 bis
		(µg)	(µg)	(µg)
Tt	1	1	_	1
Dt	1	-	3	_
Dt	3	3	3	3
Tt	. 4	1	1	1
Tt	5	1	_	1
Dt	5	_	3	
Dt	6B	10	10	3
Tt	6B	_	_	1
Tt	7F	1	_	1
Dt	7F	_	3	_
Tt	9V	1	1	1
Dt	9V	_	_	3
Dt	14	3	3	3
Dt	18C	3	3	3
Tt	18C	_	_	1
Tt	19F	1	1	1
Tt	23F	1	1	1
Dt	23F	_	_	3

The approximate protein load in each of the three formulations is as follows:

	F3	F4	F3 bis
Dt	about 40 µg	about 60 µg	about 40 µg
Tt	about 15 µg	about 8 µg	about 18 µg

(which corresponds to a protein/polysaccharide weight
10 ratio of about 2).

#### Claims

- Composition comprising "n" conjugates C1 to Cn, each conjugate being composed (i) of a polysaccharide derived from a Streptococcus pneumoniae serotype/serogroup S1 to Sn respectively, and (ii) of a carrier protein P1 to Pn respectively; "n" being a number equal to or greater than 2; in which composition the polysaccharides S1 to Sn are identical or different and
- in which the carrier proteins P1 to Pn are selected independently from a group consisting of "m" carrier proteins A1 to Am, "m" being a number equal to or greater than 2, provided that at least one of the carrier proteins P1 to Pn is different from the others.
- 2. Composition according to Claim 1, in which the conjugates C1 to Cn are all different from each other either by their polysaccharide or by their carrier protein or by their polysaccharide and their carrier protein.
- 20 3. Composition according to Claim 2, in which the polysaccharides S1 to Sn are all different from each other.
  - 4. Composition according to Claim 1, 2 or 3, in which "n" is a number equal to or greater than 6.
- 25 5. Composition according to Claim 4, in which "n" is a number equal to or greater than 10.
  - 6. Composition according to one of Claims 1 to 5, in which the carrier proteins P1 to Pn are independently selected from a group consisting of two carrier proteins A1 and A2.
  - 7. Composition according to Claim 6, in which when "n" is an even number, "n"/2 carrier proteins P1 to Pn are A1 and "n"/2 carrier proteins P1 to Pn are A2 or when "n" is an odd number, ("n"+1)/2 carrier proteins
- P1 to Pn are A1 and ("n"-1)/2 carrier proteins P1 to Pn are A2.
  - 8. Composition according to one of Claims 1 to 7, in which at least one of the carrier proteins P1 to Pn

- is the diphtheria toxoid (Dt) and at least one of the carrier proteins P1 to Pn is the tetanus toxoid (Tt).
- 9. Composition according to Claim 8, in which the carrier proteins P1 to Pn are selected from the group consisting of Dt and Tt.
- 10. Composition according to Claim 8 or 9, in which the quantity of Dt is less than or equal to  $200~\mu g/dose$ .
- 11. Composition according to Claim 8, 9 or 10, in 10 which the quantity of Tt is less than or equal to 50 µg/dose.
  - 12. Composition according to Claim 1, which comprises 10 or 11 valencies represented by 10 or 11 conjugates in which the polysaccharides are all
- different from each other and are derived from the serotypes chosen from serotypes 1, 3, 4, 5, 6B, 7F, 9V, 14, 18C, 19F and 23F of *S. pneumoniae*.
  - 13. Composition according to Claim 12, which comprises 10 or 11 conjugates selected from:
- 20 serotype 1 polysaccharide coupled to Tt or to Dt;
  - serotype 3 polysaccharide coupled to Dt;
  - serotype 4 polysaccharide coupled to Tt;
  - serotype 5 polysaccharide coupled to Tt or to Dt;
  - serotype 6B polysaccharide coupled to Dt;
- 25 serotype 7F polysaccharide coupled to Tt or to Dt;
  - serotype 9V polysaccharide coupled to Tt;
  - serotype 14 polysaccharide coupled to Dt;
  - serotype 18C polysaccharide coupled to Dt;
  - serotype 19F polysaccharide coupled to Tt; and
- 30 serotype 23F polysaccharide coupled to Tt.
  - 14. Composition according to Claim 1, comprises 10 or 11 valencies represented by 12 to 22, especially 12 to 15 conjugates. in which the polysaccharides are derived from the serotypes chosen
- from serotypes 1, 3, 4, 5, 6B, 7F, 9V, 14, 18C, 19F and 23F; in which composition conjugates of the same valency differ from each other in the carrier protein.
  - 15. Composition according to Claim 14, which comprises:

- serotype 1 polysaccharide coupled to Tt;
- serotype 3 polysaccharide coupled to Dt;
- serotype 4 polysaccharide coupled to Tt;
- serotype 5 polysaccharide coupled to Tt;
- 5 serotype 6B polysaccharide coupled to Dt;
  - serotype 6B polysaccharide coupled to Tt;
  - serotype 7F polysaccharide coupled to Tt;
  - serotype 9V polysaccharide coupled to Tt;
  - serotype 9V polysaccharide coupled to Dt;
- 10 serotype 14 polysaccharide coupled to Dt;
  - serotype 18C polysaccharide coupled to Dt;
  - serotype 18C polysaccharide coupled to Tt;
  - serotype 19F polysaccharide coupled to Tt;
  - serotype 23F polysaccharide coupled to Tt; and
- 15 serotype 23F polysaccharide coupled to Dt.
  - 16. Composition which comprises "n" conjugates C1 to Cn, each conjugate being composed (i) of a polysaccharide S1 to Sn respectively and (ii) of a carrier protein P1 to Pn respectively, "n" being a
- number equal to or greater than 2; in which composition the polysaccharides S1 to Sn are identical or different and in which the carrier proteins P1 to Pn are selected independently from a group consisting of diphtheria (Dt) and tetanus (Tt) toxoids, provided that at least
- one of the carrier proteins P1 to Pn is different from the others; and which is characterized in that the quantity of Dt and Tt is respectively less than or equal to 200 and 50 µg/dose.
- 17. Composition according to Claim 16, in which the conjugates C1 to Cn are all different from each other either by their polysaccharide or by their carrier protein or by their polysaccharide and their carrier protein.
- 18. Composition according to Claim 17, in which the polysaccharides S1 to Sn are all different from each other.
  - 19. Composition according to Claim 16, 17 or 18, in which "n" is a number equal to or greater than 6.

- 20. Composition according to Claim 19, in which "n" is a number equal to or greater than 10.
- 21. Composition according to one of Claims 16 to
- 20, in which the polysaccharides S1 to Sn are of bacterial origin.
  - 22. Composition according to Claim 21, in which the polysaccharides S1 to Sn are all derived from the same bacterial species.
- 23. Composition according to Claim 22, in which the polysaccharides S1 to Sn are all derived from the species Streptococcus pneumoniae.

Case No.: 99,849

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name.

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

#### MULTIVALENT VACCINE COMPOSITION WITH MIXED CARRIER

	, TOESTEVILLE IVI	VICERAL COM OS	THON WITH MIXED CARRIER
the specification of	which is attached !	hereto unless the follow	ring space is checked:
w	as filed on	as Unit	ed States Application Serial Number
		nd understand the conte by any amendment refe	ents of the above-identified specification, including the erred to above.
I acknowledge the d	luty to disclose inf	formation which is mate	erial to patentability as defined in 37 CFR § 1.56.
patent or inventor's country other than t	s certificate, or § the United States, int or inventor's cer	365(a) of any PCT in listed below and have a rtificate, or PCT international states and the states are states as a second state of the states are states a	19(a)-(d) or § 365(b) of any foreign application(s) for ternational application which designated at least one also identified below, by checking the box, any foreign tional application having a filing date before that of the
Prior Foreign Appli	cation(s):		
Number 1. PCT/FR98, 2. 97/06210	/00966 <b>~</b> PC	Country CT rance	Day/Month/Year Filed 05 May 1998 14 May 1997
I hereby claim the b Application 1. 2.		J.S.C. § 119(e) of any Un <u>Filing Date</u>	nited States provisional application(s) listed below:
international applications of this application provided by the first to patentability as	ation designating to cation is not disclost t paragraph of 35 to defined in 37 C.	the United States, listed osed in the prior United U.S.C. § 112, I acknowl	United States application(s), or § 365(c) of any PCT below and, insofar as the subject matter of each of the States or PCT international application in the manner ledge the duty to disclose information which is materia came available between the filing date of the prior of this application.

Status: patented, pending, abandoned

Filing Date

Application Number

I hereby appoint the following attorneys and agent(s) to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith:

Denis A. Berntsen John J. McDonnell Daniel A. Boehnen Bradley J. Hulbert Paul H. Berghoff Grantland G. Drutchas Steven J. Sarussi David M. Frischkorn James C. Gumina A. Blair Hughes Thomas A. Fairhall Emily Miao	Reg. No. 26707 Reg. No. 26949 Reg. No. 28399 Reg. No. 30130 Reg. No. 30243 Reg. No. 32565 Reg. No. 32784 Reg. No. 32833 Reg. No. 32898 Reg. No. 32901 Reg. No. 34591	Anthoula Pomrening (agent) George I. Lee James M McCarthy Jeremy Noe (agent) Sean M. Sullivan Amir N. Penn Patrick J. Halloran (agent) Joshua R. Rich Thomas E. Wettermann Vernon W. Francissen Robert J. Irvine	Reg. No. 38805 Reg. No. 39269 Reg. No. 39296 Reg. No. 40104 Reg. No. 40191 Reg. No. 40767 Reg. No. 41053 Reg. No. 41269 Reg. No. 41523 Reg. No. 41762 Reg. No. 41865
Kevin E. Noonan	Reg. No. 35285 Reg. No. 35303	Richard A. Machonkin David S. Harper	Reg. No. 41962 Reg. No. 42636
Leif R. Sigmond, Jr.	Reg. No. 35680	Stephen Lesavich	Reg. No. 43749
Lawrence H. Aaronson	Reg. No. 35818	Enrique Perez	Reg. No. 43853
Matthew J. Sampson	Reg. No. 35999	Marcus J. Thymian	Reg. No. 43954
Curt J. Whitenack	Reg. No. 36054	S. Richard Carden (agent)	Reg. No. 44588
Christopher M. Cavan	Reg. No. 36475	Mark Chael (agent)	Reg. No. 44601
Michael S. Greenfield	Reg. No. 37142	Stephen H. Docter	Reg. No. 44659
Roger P. Zimmerman	Reg. No. 38670		

Address all telephone calls to Michael S. Greenfield at (312) 913-0001. Address all correspondence to MCDONNELL BOEHNEN HULBERT & BERGHOFF, 300 South Wacker Drive, Chicago, Illinois 60606 USA.

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Full name of first inventor: Odile Leroy

Inventor's signature: Residence: France

Citizenship: France -

Post Office Address: 13, rue des Quatre Vents, F-92360 Garches F